Allergy Action Plan Place Student's D.O.B: ____Teacher:____ Name:___ Child's ALLERGY TO: Picture Here Asthmatic Yes* | No | *Higher risk for severe reaction ◆ STEP 1: TREATMENT ◆ Give Checked Medication**: Symptoms: **(To be determined by physician authorizing treatment) ☐ Antihistamine ☐ Epinephrine If a food allergen has been ingested, but no symptoms: ☐ Antihistamine □ Epinephrine Itching, tingling, or swelling of lips, tongue, mouth Mouth ☐ Antihistamine ☐ Epinephrine Hives, itchy rash, swelling of the face or extremities Skin ☐ Epinephrine ☐ Antihistamine Nausea, abdominal cramps, vomiting, diarrhea 8 Gut ☐ Epinephrine ☐ Antihistamine Tightening of throat, hoarseness, hacking cough Throat† ☐ Antihistamine ☐ Epinephrine Shortness of breath, repetitive coughing, wheezing Lung† ☐ Antihistamine ☐ Epinephrine Weak or thready pulse, low blood pressure, fainting, pale, blueness Heart† 框 ☐ Antihistamine ☐ Epinephrine Other† ☐ Epinephrine ☐ Antihistamine If reaction is progressing (several of the above areas affected), give: †Potentially life-threatening. The severity of symptoms can quickly change. DOSAGE Epinephrine: inject intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject® 0.3 mg Twinject® 0.15 mg (see reverse side for instructions) Antihistamine: give_____ medication/dose/route Other: give medication/dose/route IMPORTANT: Asthma inhalers and/or antihistamines cannot be depended on to replace epinephrine in anaphylaxis. ◆ STEP 2: EMERGENCY CALLS ◆ 1. Call 911 (or Rescue Squad: ______). State that an allergic reaction has been treated, and additional epinephrine may be needed. Phone Number: 2. Dr. _____ Phone Number(s) 3. Parent_____ 4. Emergency contacts: Phone Number(s) Name/Relationship 2.) 1.) EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE CHILD TO MEDICAL FACILITY! Parent/Guardian's Signature_____

Doctor's Signature_____

(Required)